




Operation Information

Catheter Ablation of Atrial Fibrillation

Introduction

Atrial Fibrillation (AF) is an irregular and chaotic heart rhythm that occurs in the upper chambers of the heart (atria). It may cause impairment of contraction of the atrium, and formation of blood clots. Although AF is not life-threatening, individuals may experience symptoms such as palpation, shortness of breath, chest discomfort, fatigue and fainting. Older clients have a higher chance of developing a stroke if they have AF.

Catheter ablation of atrial fibrillation is an operation on the heart that aims to resolve AF. The doctor uses specially designed needles and instruments to perform transseptal left heart catheterization by making a hole in the area separating the right and left atrium, in order to pass the catheters from the right atrium to the left atrium. Energy is released from the catheter, and encircles the entrance of the veins. The ablation changes the heart muscle inside the atrium near the pulmonary veins to block the abnormal signals from entering the heart.



Source:
<https://consultqd.clevelandclinic.org/ablation-vs-medical-therapy-for-atrial-fibrillation-putting-cabana-in-perspective/>

Outcomes

This operation cures AF, relieves symptoms and improve the individual's quality of life. Please discuss with your doctor to select the most appropriate treatment.

Procedures

1. This operation can be performed under local anaesthesia, general anaesthesia or monitored anaesthetic care (MAC) in a cardiac catheterization laboratory. If local anaesthesia is used, the doctor may give you sedation to calm you down.
2. Electrodes are adhered to your chest to monitor the heart rate and rhythm. Blood oxygen and blood pressure are monitored through the devices on your fingertip and arm.
3. Small wounds are made over the groin, under the clavicle or around the neck for the access to arteries or veins.
4. Catheters are advanced to the heart under X-ray guidance.
5. Since the abnormal electrical activities usually arise from the left atrium, the doctor needs to perform transseptal left heart catheterization, to allow the passage of catheters from the right atrium to the left atrium.
6. At specific sites inside the heart, the doctor records electrical information, then a tiny electric current is delivered to alter the heart impulse and try to trigger arrhythmia.
7. You may experience discomfort when the heart is being excited to a certain rate. When an induced arrhythmia persists, the doctor may use direct current cardioversion to convert it.
8. Energy is delivered to the target site via a special catheter. You may experience some chest discomfort during the delivery of energy.
9. After removing all the catheters, pressure will be applied to the puncture site to stop bleeding.
10. You are sent to a ward for close observation for 12 to 24 hours.

Possible Risks and Complications

Minor complications:

- Wound infection
- Wound bleeding

Major complications (4.5%):

- Damage of blood vessels, lungs and heart that may need surgical intervention
- Cardiac tamponade
- Narrowing of the pulmonary vein
- Damage of the nerve supplying the diaphragm which leads to permanent diaphragmatic paralysis
- Stroke
- Formation of abnormal communication between the esophagus and the heart
- Infection
- Death due to uncontrollable complications (0.15%)

Other complications:

- May not be able to improve symptoms in about 30% of cases
- Some clients may require more than one catheter ablation

** The risks listed above are in general terms and the possibility of complications is not exhaustive. Please understand that even though all procedures are carried out with utmost professionalism and care, this does not rule out the possibility of complications from arising. In the event of peripheral organ damage or post-operative haemorrhage or leakage, further operations may be required.

Pre-operative Preparations

1. Good hygiene can prevent surgical wound infection. Therefore, we advise you to clean up yourself on the day of operation.
2. The procedure and possible complications will be explained by the doctor and a consent form must be signed prior to the operation.
3. Please inform the doctor and nurse all your past medical history, previous surgical operations, current medication and any complication with drug or anaesthesia.
4. If you are a female, please provide the date of your last menstrual period (LMP) and avoid pregnancy before the operation. This operation involves radiation exposure.
5. You may be instructed to discontinue some or all of the anti-arrhythmic drugs and blood thinners (e.g. Warfarin, Xarelto or Pradaxa) before the operation.
6. You may need to undergo some investigations like blood tests, electrocardiogram (ECG), chest X-ray, computerized tomography (CT) or magnetic resonance imaging (MRI) of the heart and transesophageal echocardiogram to ensure there is no abnormal blood clot and rule out other structural heart diseases.
7. Shaving may be required on the puncture site(s).
8. No food or drink four to six hours before the operation, or situation dependent.
9. Please change into a surgical gown after removing all belongings including undergarments, dentures, jewellery and contact lenses.
10. Please empty your bladder before the operation.

Post-operative Instructions

1. Vital signs and the wound are monitored closely by the nursing staff.
2. Please inform the nurse immediately if pain or bleeding from the puncture site, numbness of the affected limb, difficulty breathing or any other discomfort.
3. You are required to bed rest for at least 4 hours. Please do not move or bend the affected limb. Whenever you cough or sneeze, please apply pressure on the puncture site with your hand.

Advice on Discharge

1. The wound is covered with a waterproof dressing. Dressing can be changed on the day after the operation. Always keep the wound site clean and dry.
2. Showers are allowed with waterproof dressing if no abnormality is found on the wound. Be reminded to pat the puncture site and surrounding area dry and change a new waterproof dressing as instructed afterward.
3. Bruising or mild swelling around the puncture site is common and usually subsides 2-3 weeks later.
4. Avoid any lifting or strenuous activity for one week, to prevent increasing pressure which may cause the puncture site re-bleeding.
5. It is common to experience palpitations on and off for the first 3 months after the operation, thus you still need to continue anti-arrhythmic medication. The final result of the operation will be more obvious after 3 months.
6. No driving for 1 week. Please strictly follow the doctor's instructions.
7. You may need to take blood thinner to prevent the formation and dislodgment of blood clots from the heart for a certain period of time. Depending on the risk of future stroke, your doctor will determine whether you need long-term blood thinner.
8. It is rare to have severe bleeding from the puncture site. If bleeding does occur you must lie flat with the affected limb elevated, and apply pressure to the site for at least 10 minutes. For unstoppped or severe bleeding, please call 999 for emergency help.
9. Immediately consult your doctor or return to hospital for professional attention in the event of bleeding, pus discharge, increasing pain and swelling from the wound, chest pain, breathing difficulty, shivering, high fever over 38°C or 100.4°F, or any other unusual symptoms.
10. Any follow-up consultations should be attended as scheduled.

Should there be any enquiries or concerns, please consult the attending doctor.

Under the professional care of the doctor, you will gradually recover. We wish you all the best during your treatment and recovery.

If you have any questions after reading the entire leaflet, please write them down in the spaces provided in order for the doctor to further follow-up.

Compiled by Union Hospital Operating Theatre (OT) Governance Committee

The above information is for reference only, please enquire your physician for details
Our Hospital reserves the RIGHT to amend any information in this leaflet without prior notification